



Hamdy O, Carver C: The Why WAIT program: improving clinical outcomes through weight management in type 2 diabetes, *Curr Diabetes Rep*. 8:413–420, 2008.

Ikramuddin S, Korner J, Lee WJ, et al: Roux-en-Y gastric bypass vs intensive medical management for the control of type 2 diabetes, hypertension, and hyperlipidemia: the Diabetes Surgery Study randomized clinical trial, *JAMA* 309:2240–2249, 2013.

Look AHEAD Research Group, Wing RR, Bolin P, et al: Cardiovascular effects of intensive lifestyle intervention in type 2 diabetes, *N Engl J Med* 369:145–154, 2013.

Maggard MA, Shugarman LR, Suttorp M, et al: Meta-analysis: surgical treatment of obesity, *Ann Intern Med* 142:547–559, 2005.

Schaer PR, Kashyap SR, Wolski K, et al: Bariatric surgery versus intensive medical therapy in obese patients with diabetes, *N Engl J Med* 366:1567–1576, 2012.

Sjostrom L, Lindroos AK, Peltonen M, et al: Swedish Obese Subjects Study Scientific Group. Lifestyle, diabetes, and cardiovascular risk factors 10 years after bariatric surgery, *N Engl J Med* 351:2683–2693, 2004.

Strauss RS, Bradley LJ, Brolin RE: Gastric bypass surgery in adolescents with morbid obesity, *J Pediatr* 138:499–504, 2001.