

children (i.e., proxy). All of these individuals require objective but complete, tactful, and compassionate assessments and considerable psychiatric care.

#### SUGGESTED READINGS

- Aduan RP, Fauci AS, Dale DC, et al: Prolonged fever of unknown origin (FUO): a prospective study of 347 patients, *Clin Res* 26:558A, 1978.
- Aduan RP, Fauci AS, Dale DC, et al: Factitious fever and self-induced infection: a report of 32 cases and review of the literature, *Ann Intern Med* 90:230–242, 1979.
- Cannon J: Perspective on fever: the basic science and conventional medicine, *Complement Ther Med* 21(Suppl 1):S54–S60, 2013.
- Hayakawa K, Balaji R, Pranatharthi C: Fever of unknown origin: an evidence-based review, *Am J Med Sci* 344:307–316, 2012.
- Rezai-Zadeh K, Munzberg H: Integration of sensory information via central thermoregulatory leptin targets, *Physiol Behav* 121:49–55, 2013.
- Weber D, Cohen M, Morrell D: The acutely ill patient with fever and rash. In Mandell GL, Bennett JE, Dolin R, editors: *Mandell, Douglas, and Bennett's principles and practice of infectious diseases*, ed 7, Philadelphia, 2010, Churchill Livingstone, pp 791–807.

