

criminating chronotropic incompetence from resting bradycardia and may aid in the identification of the mechanism of exercise intolerance.

Autonomic nervous system testing is useful in diagnosing carotid sinus hypersensitivity; pauses >3 s are consistent with the diagnosis but may be present in asymptomatic elderly subjects. Determining the intrinsic heart rate (IHR) may distinguish SA node dysfunction from slow heart rates that result from high vagal tone. The normal IHR after administration of 0.2 mg/kg propranolol and 0.04 mg/kg atropine is $117.2 - (0.53 \times \text{age})$ in beats/min; a low IHR is indicative of SA disease.

Electrophysiologic testing may play a role in the assessment of patients with presumed SA node dysfunction and in the evaluation of syncope, particularly in the setting of structural heart disease. In this circumstance, electrophysiologic testing is used to rule out more malignant etiologies of syncope, such as ventricular tachyarrhythmias and AV conduction block. There are several ways to assess SA node function invasively. They include the sinus node recovery time (SNRT), defined as the longest pause after cessation of overdrive pacing of the right atrium near the SA node (normal: <1500 ms or, corrected for sinus cycle length, <550 ms), and the sinoatrial conduction time (SACT), defined as one-half the difference between the intrinsic sinus cycle length and a noncompensatory pause after a premature atrial stimulus (normal <125 ms). The combination of an abnormal SNRT, an abnormal SACT, and a low IHR is a sensitive and specific indicator of intrinsic SA node disease.

TREATMENT SINOATRIAL NODE DYSFUNCTION

Since SA node dysfunction is not associated with increased mortality rates, the aim of therapy is alleviation of symptoms. Exclusion of extrinsic causes of SA node dysfunction and correlation of the cardiac rhythm with symptoms is an essential part of patient management. Pacemaker implantation is the primary therapeutic intervention in patients with symptomatic SA node dysfunction. Pharmacologic considerations are important in the evaluation and management of patients with SA nodal disease. A number of drugs modulate SA node function and are extrinsic causes of dysfunction (Table 274-1). Beta blockers and calcium channel blockers increase SNRT in patients with SA node dysfunction, and antiarrhythmic drugs with class I and III action may promote SA node exit block. In general, such agents should be discontinued before decisions regarding the need for permanent pacing in patients with SA node disease are made. Chronic pharmacologic therapy for sinus bradyarrhythmias is limited. Some pharmacologic agents may improve SA node function; digitalis, for example, has been shown to shorten SNRT in patients with SA node dysfunction. Isoproterenol or atropine administered IV may increase the sinus rate acutely. Theophylline has been used both acutely and chronically to increase heart rate but has liabilities when used in patients with tachycardia-bradycardia syndrome, increasing the frequency of supraventricular tachyarrhythmias, and in patients with structural heart disease, increasing the risk of potentially serious ventricular arrhythmias. Currently, there is only a single randomized study of therapy for SA node dysfunction. In patients with resting heart rates <50 and >30 beats/min on a Holter monitor, patients who received dual-chamber pacemakers experienced significantly fewer syncopal episodes and had symptomatic improvement compared with patients randomized to theophylline or no treatment.

In certain circumstances, sinus bradycardia requires no specific treatment or only temporary rate support. Sinus bradycardia is common in patients with acute inferior or posterior MI and can be exacerbated by vagal activation induced by pain or the use of drugs such as morphine. Ischemia of the SA nodal artery probably occurs in acute coronary syndromes more typically with involvement with the right coronary artery, and even with infarction, the effect on SA node function most often is transient.

Sinus bradycardia is a prominent feature of carotid sinus hypersensitivity and neurally mediated hypotension associated with vasovagal syncope that responds to pacemaker therapy. Carotid

hypersensitivity with recurrent syncope or presyncope associated with a predominant cardioinhibitory component responds to pacemaker implantation. Several randomized trials have investigated the efficacy of permanent pacing in patients with drug-refractory vasovagal syncope, with mixed results. Although initial trials suggested that patients undergoing pacemaker implantation have fewer recurrences and a longer time to recurrence of symptoms, at least one follow-up study did not confirm these results.

PERMANENT PACEMAKERS

Nomenclature and Complications The main therapeutic intervention in SA node dysfunction is permanent pacing. Since the first implementation of permanent pacing in the 1950s, many advances in technology have resulted in miniaturization, increased longevity of pulse generators, improvement in leads, and increased functionality. To better understand pacemaker therapy for bradycardias, it is important to be familiar with the fundamentals of pacemaking. Pacemaker modes and function are named using a five-letter code. The first letter indicates the chamber(s) that is paced (O, none; A, atrium; V, ventricle; D, dual; S, single), the second is the chamber(s) in which sensing occurs (O, none; A, atrium; V, ventricle; D, dual; S, single), the third is the response to a sensed event (O, none; I, inhibition; T, triggered; D, inhibition + triggered), the fourth refers to the programmability or rate response (R, rate responsive), and the fifth refers to the existence of antitachycardia functions if present (O, none; P, antitachycardia pacing; S, shock; D, pace + shock). Almost all modern pacemakers are multiprogrammable and have the capability for rate responsiveness using one of several rate sensors: activity or motion, minute ventilation, or QT interval. The most commonly programmed modes of implanted single- and dual-chamber pacemakers are VVIR and DDDR, respectively, although multiple modes can be programmed in modern pacemakers.

Although pacemakers are highly reliable, they are subject to a number of complications related to implantation and electronic function. In adults, permanent pacemakers are most commonly implanted with access to the heart by way of the subclavian–superior vena cava venous system. Rare, but possible, acute complications of transvenous pacemaker implantation include infection, hematoma, pneumothorax, cardiac perforation, diaphragmatic/phrenic nerve stimulation, and lead dislodgment. Limitations of chronic pacemaker therapy include infection, erosion, lead failure, and abnormalities resulting from inappropriate programming or interaction with the patient's native electrical cardiac function. Rotation of the pacemaker pulse generator in its subcutaneous pocket, either intentionally or inadvertently, often referred to as "twiddler's syndrome," can wrap the leads around the generator and produce dislodgment with failure to sense or pace the heart. The small size and light weight of contemporary pacemakers make this a rare complication.

Complications stemming from chronic cardiac pacing also result from disturbances in atrioventricular synchrony and/or left ventricular mechanical synchrony. Pacing modes that interrupt or fail to restore atrioventricular synchrony may lead to a constellation of signs and symptoms, collectively referred to as pacemaker syndrome, that include neck pulsation, fatigue, palpitations, cough, confusion, exertional dyspnea, dizziness, syncope, elevation in jugular venous pressure, canon A waves, and stigmata of congestive heart failure, including edema, rales, and a third heart sound. Right ventricular apical pacing can induce dyssynchronous activation of the left ventricle, leading to compromised left ventricular systolic function, mitral valve regurgitation, and the previously mentioned stigmata of congestive heart failure. Maintenance of AV synchrony can minimize the sequelae of pacemaker syndrome. Selection of pacing modes that minimize unnecessary ventricular pacing or implantation of a device capable of right and left ventricular pacing (biventricular pacing) can help minimize the deleterious consequences of pacing-induced mechanical dyssynchrony at the ventricular level.