



FIGURE 421-1 The density and size distribution of the major classes of lipoprotein particles. Lipoproteins are classified by density and size, which are inversely related. HDL, high-density lipoprotein; IDL, intermediate-density lipoprotein; LDL, low-density lipoprotein; VLDL, very-low-density lipoprotein.

(plasma, interstitial fluid, and lymph) to and from tissues. Lipoproteins play an essential role in the absorption of dietary cholesterol, long-chain fatty acids, and fat-soluble vitamins; the transport of triglycerides, cholesterol, and fat-soluble vitamins from the liver to peripheral tissues; and the transport of cholesterol from peripheral tissues to the liver and intestine.

Lipoproteins contain a core of hydrophobic lipids (triglycerides and cholesteryl esters) surrounded by a shell of hydrophilic lipids (phospholipids, unesterified cholesterol) and proteins (called apolipoproteins) that interact with body fluids. The plasma lipoproteins are divided into five major classes based on their relative density (Fig. 421-1 and Table 421-1): chylomicrons, very-low-density lipoproteins (VLDLs), intermediate-density lipoproteins (IDLs), low-density lipoproteins (LDLs), and high-density lipoproteins (HDLs). Each lipoprotein class comprises a family of particles that vary in density, size, and protein composition. Because lipid is less dense than water, the density of a lipoprotein particle is primarily determined by the amount of lipid per particle. Chylomicrons are the most lipid-rich and therefore least dense lipoprotein particles, whereas HDLs have the least lipid and are therefore the most dense lipoproteins. In addition to their density, lipoprotein particles can be classified according to their size, determined either by non-denaturing gel electrophoresis

TABLE 421-2 MAJOR APOLIPOPROTEINS

Apolipoprotein	Primary Source	Lipoprotein Association	Function
ApoA-I	Intestine, liver	HDL, chylomicrons	Structural protein for HDL Activates LCAT
ApoA-II	Liver	HDL, chylomicrons	Structural protein for HDL
ApoA-IV	Intestine, liver	HDL, chylomicrons	Unknown
ApoA-V	Liver	VLDL, chylomicrons	Promotes LPL-mediated triglyceride lipolysis
Apo(a)	Liver	Lp(a)	Unknown
ApoB-48	Intestine	Chylomicrons, chylomicron remnants	Structural protein for chylomicrons
ApoB-100	Liver	VLDL, IDL, LDL, Lp(a)	Structural protein for VLDL, LDL, IDL, Lp(a) Ligand for binding to LDL receptor
ApoC-I	Liver	Chylomicrons, VLDL, HDL	Unknown
ApoC-II	Liver	Chylomicrons, VLDL, HDL	Cofactor for LPL
ApoC-III	Liver, intestine	Chylomicrons, VLDL, HDL	Inhibits LPL activity and lipoprotein binding to receptors
ApoE	Liver	Chylomicron remnants, IDL, HDL	Ligand for binding to LDL receptor and other receptors

Abbreviations: HDL, high-density lipoprotein; IDL, intermediate-density lipoprotein; LCAT, lecithin-cholesterol acyltransferase; LDL, low-density lipoprotein; Lp(a), lipoprotein A; LPL, lipoprotein lipase; VLDL, very-low-density lipoprotein.

or by nuclear magnetic resonance profiling. There is a strong inverse relationship between density and size, with the largest particles being the most buoyant (chylomicrons) and the smallest particles being the most dense (HDL).

The proteins associated with lipoproteins, called *apolipoproteins* (Table 421-2), are required for the assembly, structure, function, and metabolism of lipoproteins. Apolipoproteins activate enzymes important in lipoprotein metabolism and act as ligands for cell surface receptors. ApoB is a very large protein and is the major structural protein of chylomicrons, VLDLs, IDLs, and LDLs; one molecule

TABLE 421-1 MAJOR LIPOPROTEIN CLASSES

Lipoprotein	Density, g/mL ^a	Size, nm ^b	Electrophoretic Mobility ^c	Apolipoproteins		
				Major	Other	Other Constituents
Chylomicrons	0.930	75–1200	Origin	ApoB-48	A-I, A-V, C-I, C-II, C-III, E	Retinyl esters
Chylomicron remnants	0.930–1.006	30–80	Slow pre-β	ApoB-48	A-I, A-V, C-I, C-II, C-III, E	Retinyl esters
VLDL	0.930–1.006	30–80	Pre-β	ApoB-100	A-I, A-II, A-V, C-I, C-II, C-III, E	Vitamin E
IDL	1.006–1.019	25–35	Slow pre-β	ApoB-100	C-I, C-II, C-III, E	Vitamin E
LDL	1.019–1.063	18–25	β	ApoB-100		Vitamin E
HDL	1.063–1.210	5–12	α	ApoA-I	A-II, A-IV, A-V, C-III, E	LCAT, CETP, paroxonase
Lp(a)	1.050–1.120	25	Pre-β	ApoB-100	Apo(a)	Oxidized phospholipids

^aThe density of the particle is determined by ultracentrifugation. ^bThe size of the particle is measured using gel electrophoresis. ^cThe electrophoretic mobility of the particle on agarose gel electrophoresis reflects the size and surface charge of the particle, with β being the position of LDL and α being the position of HDL.

Note: All of the lipoprotein classes contain phospholipids, esterified and unesterified cholesterol, and triglycerides to varying degrees.

Abbreviations: CETP, cholesteryl ester transfer protein; HDL, high-density lipoprotein; IDL, intermediate-density lipoprotein; LCAT, lecithin-cholesterol acyltransferase; LDL, low-density lipoprotein; Lp(a), lipoprotein A; VLDL, very-low-density lipoprotein.