

TABLE 43-1 HEREDITARY HEARING IMPAIRMENT GENES

Designation	Gene	Function	Designation	Gene	Function
Autosomal Dominant			DFNB12	<i>CDH23</i>	Intercellular adherence protein
DFNA1	<i>CRYM</i>	Thyroid hormone-binding protein	DFNB15/72/95	<i>GIPC3</i>	PDZ domain containing protein
DFNA2A	<i>DIAPH1</i>	Cytoskeletal protein	DFNB16	<i>STRC</i>	Stereocilia protein
DFNA2B	<i>KCNQ4</i>	Potassium channel	DFNB18	<i>USH1C</i>	Unknown
DFNA3A	<i>GJB3 (Cx31)</i>	Gap junction	DFNB21	<i>TECTA</i>	Tectorial membrane protein
DFNA3B	<i>GJB2 (Cx26)</i>	Gap junction	DFNB22	<i>OTOA</i>	Gel attachment to nonsensory cell
DFNA4	<i>GJB6 (Cx30)</i>	Gap junction	DFNB23	<i>PCDH15</i>	Morphogenesis and cohesion
	<i>MYH14</i>	Class II nonmuscle myosin	DFNB24	<i>RDX</i>	Cytoskeletal protein
	<i>CEACAM16</i>	Cell adhesion molecule	DFNB25	<i>GRXCR1</i>	Reversible S-glutathionylation of proteins
DFNA5	<i>DFNA5</i>	Unknown	DFNB28	<i>TRIOBP</i>	Cytoskeletal-organizing protein
DFNA6/14/38	<i>WFS1</i>	Transmembrane protein	DFNB29	<i>CLDN14</i>	Tight junctions
DFNA8/12	<i>TECTA</i>	Tectorial membrane protein	DFNB30	<i>MYO3A</i>	Hybrid motor-signaling myosin
DFNA9	<i>COCH</i>	Unknown	DFNB31	<i>WHRN</i>	PDZ domain-containing protein
DFNA10	<i>EYA4</i>	Developmental gene	DFNB35	<i>ESRRB</i>	Estrogen-related receptor beta protein
DFNA11	<i>MYO7A</i>	Cytoskeletal protein	DFNB36	<i>ESPN</i>	Ca-insensitive actin-bundling protein
DFNA13	<i>COL11A2</i>	Cytoskeletal protein	DFNB37	<i>MYO6</i>	Unconventional myosin
DFNA15	<i>POU4F3</i>	Transcription factor	DFNB39	<i>HFG</i>	Hepatocyte growth factor
DFNA17	<i>MYH9</i>	Cytoskeletal protein	DFNB42	<i>ILDR1</i>	Ig-like domain-containing receptor
DFNA20/26	<i>ACTG1</i>	Cytoskeletal protein	DFNB48	<i>CIB2</i>	Calcium and integrin binding protein
DFNA22	<i>MYO6</i>	Unconventional myosin	DFNB49	<i>MARVELD2</i>	Tight junction protein
DFNA23	<i>SIX1</i>	Developmental gene	DFNB53	<i>COL11A2</i>	Collagen protein
DFNA25	<i>SLC17A8</i>	Vesicular glutamate transporter	DFNB59	<i>PJVK</i>	Zn-binding protein
DFNA28	<i>TFCP2L3</i>	Transcription factor	DFNB61	<i>SLC26A5</i>	Motor protein
DFNA36	<i>TMC1</i>	Transmembrane protein	DFNB63	<i>LRTOMT/COMT2</i>	Putative methyltransferase
DFNA41	<i>P2RX2</i>	Purinergic receptor	DFNB66/67	<i>LHFPL5</i>	Tetraspan protein
DFNA44	<i>CCDC50</i>	Effector of epidermal growth factor-mediated signaling	DFNB70	<i>PNPT1</i>	Mitochondrial-RNA-import protein
DFNA48	<i>MYO1A</i>	Unconventional myosin	DFNB74	<i>MSRB3</i>	Methionine sulfoxide reductase
DFNA50	<i>MIRN96</i>	MicroRNA	DFNB77	<i>LOXHD1</i>	Stereociliary protein
DFNA51	<i>TJP2</i>	Tight junction protein	DFNB79	<i>TPRN</i>	Unknown
DFNA56	<i>TNC</i>	Extracellular matrix protein	DFNB82	<i>GPSM2</i>	G protein signaling modulator
DFNA64	<i>SMAC/DIABLO</i>	Mitochondrial proapoptotic protein	DFNB84	<i>PTPRQ</i>	Type III receptor-like protein-tyrosine phosphatase family
Autosomal Recessive			DFNB86	<i>TBC1D24</i>	GTPase-activating protein
DFNB1A	<i>GJB2 (Cx26)</i>	Gap junction	DFNB88	<i>ELMOD3</i>	GTPase-activating protein
DFNB1B	<i>GJB6 (Cx30)</i>	Gap junction	DFNB89	<i>KARS</i>	Lysyl-tRNA synthetase
DFNB2	<i>MYO7A</i>	Cytoskeletal protein	DFNB91	<i>GJB3</i>	Gap junction
DFNB3	<i>MYO15</i>	Cytoskeletal protein	DFNB93	<i>CABP2</i>	Calcium binding protein
DFNB4	<i>PDS (SLC26A4)</i>	Chloride/iodide transporter	DFNB98	<i>TSPEAR</i>	Epilepsy-associated repeats containing protein
DFNB6	<i>TMIE</i>	Transmembrane protein		<i>SERPIN6</i>	Protease inhibitor
DFNB7/B11	<i>TMC1</i>	Transmembrane protein			
DFNB9	<i>OTOF</i>	Trafficking of membrane vesicles			
DFNB8/10	<i>TMPRSS3</i>	Transmembrane serine protease			

The Rinne and Weber tuning fork tests, with a 512-Hz tuning fork, are used to screen for hearing loss, differentiate conductive from sensorineural hearing losses, and confirm the findings of audiologic evaluation. The Rinne test compares the ability to hear by air conduction with the ability to hear by bone conduction. The tines of a vibrating tuning fork are held near the opening of the external auditory canal, and then the stem is placed on the mastoid process; for direct contact, it may be placed on teeth or dentures. The patient is asked to indicate whether the tone is louder by air conduction or bone conduction. Normally, and in the presence of sensorineural hearing loss, a tone is heard louder by air conduction than by bone conduction; however, with conductive hearing loss of ≥ 30 dB (see "Audiologic Assessment," below), the bone-conduction stimulus is perceived as louder than the air-conduction stimulus. For the Weber test, the stem of a vibrating tuning fork is placed on

the head in the midline and the patient is asked whether the tone is heard in both ears or better in one ear than in the other. With a unilateral conductive hearing loss, the tone is perceived in the affected ear. With a unilateral sensorineural hearing loss, the tone is perceived in the unaffected ear. A 5-dB difference in hearing between the two ears is required for lateralization.

LABORATORY ASSESSMENT OF HEARING

Audiologic Assessment The minimum audiologic assessment for hearing loss should include the measurement of pure tone air-conduction and bone-conduction thresholds, speech reception threshold, word recognition score, tympanometry, acoustic reflexes, and acoustic-reflex decay. This test battery provides a screening evaluation of the entire auditory system and allows one to determine whether further