

**TABLE 77-5** NORMAL MARROW RESPONSE TO ANEMIA


Hemoglobin	Production Index	Reticulocyte Count
15 g/dL	1	50,000/ $\mu$ L
11 g/dL	2.0–2.5	100–150,000/ $\mu$ L
8 g/dL	3.0–4.0	300–400,000/ $\mu$ L

corrected reticulocyte count is the *reticulocyte production index*, and it provides an estimate of marrow production relative to normal. In many hospital laboratories, the reticulocyte count is reported not only as a percentage but also in absolute numbers. If so, no correction for dilution is required. A summary of the appropriate marrow response to varying degrees of anemia is shown in **Table 77-5**.


Premature release of reticulocytes is normally due to increased EPO stimulation. However, if the integrity of the bone marrow release process is lost through tumor infiltration, fibrosis, or other disorders, the appearance of nucleated red cells or polychromatophilic macrocytes should still invoke the second reticulocyte correction. The shift correction should always be applied to a patient with anemia and a very high reticulocyte count to provide a true index of effective red cell production. Patients with severe chronic hemolytic anemia may increase red cell production as much as six- to sevenfold. This measure alone confirms the fact that the patient has an appropriate EPO response, a normally functioning bone marrow, and sufficient iron available to meet the demands for new red cell formation. If the reticulocyte production index is  $<2$  in the face of established anemia, a defect in erythroid marrow proliferation or maturation must be present.

**Tests of Iron Supply and Storage** The laboratory measurements that reflect the availability of iron for hemoglobin synthesis include the serum iron, the TIBC, and the percent transferrin saturation. The percent transferrin saturation is derived by dividing the serum iron level ( $\times 100$ ) by the TIBC. The normal serum iron ranges from 9 to 27  $\mu$ mol/L (50–150  $\mu$ g/dL), whereas the normal TIBC is 54–64  $\mu$ mol/L (300–360  $\mu$ g/dL); the normal transferrin saturation ranges from 25 to 50%. A diurnal variation in the serum iron leads to a variation in the percent transferrin saturation. The serum ferritin is used to evaluate total body iron stores. Adult males have serum ferritin levels that average  $\sim 100$   $\mu$ g/L, corresponding to iron stores of  $\sim 1$  g. Adult females have lower serum ferritin levels averaging 30  $\mu$ g/L, reflecting lower iron stores ( $\sim 300$  mg). A serum ferritin level of 10–15  $\mu$ g/L indicates depletion of body iron stores. However, ferritin is also an acute-phase reactant and, in the presence of acute or chronic inflammation, may rise several-fold above baseline levels. As a rule, a serum ferritin  $>200$   $\mu$ g/L means there is at least some iron in tissue stores.


**Bone Marrow Examination** A bone marrow aspirate and smear or a needle biopsy can be useful in the evaluation of some patients with anemia. In patients with hypoproliferative anemia and normal iron status, a bone marrow is indicated. Marrow examination can diagnose primary marrow disorders such as myelofibrosis, a red cell maturation defect, or an infiltrative disease (**Figs. 77-14 to 77-16**). The increase or decrease of one cell lineage (myeloid vs erythroid) compared to another is obtained by a differential count of nucleated cells in a bone marrow smear (the myeloid/erythroid [M/E] ratio). A patient with a hypoproliferative anemia (see below) and a reticulocyte production index  $<2$  will demonstrate an M/E ratio of 2 or 3:1. In contrast, patients with hemolytic disease and a production index  $>3$  will have an M/E ratio of at least 1:1. Maturation disorders are identified from the discrepancy between the M/E ratio and the reticulocyte production index (see below). Either the marrow smear or biopsy can be stained for the presence of iron stores or iron in developing red cells. The storage iron is in the form of ferritin or *hemosiderin*. On carefully prepared bone marrow smears, small ferritin granules can normally be seen under oil immersion in 20–40% of developing erythroblasts. Such cells are called *sideroblasts*.



**FIGURE 77-14** Normal bone marrow. This is a low-power view of a section of a normal bone marrow biopsy stained with hematoxylin and eosin (H&E). Note that the nucleated cellular elements account for  $\sim 40$ – $50\%$  and the fat (clear areas) accounts for  $\sim 50$ – $60\%$  of the area. (From RS Hillman et al: *Hematology in Clinical Practice*, 5th ed. New York, McGraw-Hill, 2010.)



**FIGURE 77-15** Erythroid hyperplasia. This marrow shows an increase in the fraction of cells in the erythroid lineage as might be seen when a normal marrow compensates for acute blood loss or hemolysis. The myeloid/erythroid (M/E) ratio is about 1:1. (From RS Hillman et al: *Hematology in Clinical Practice*, 5th ed. New York, McGraw-Hill, 2010.)



**FIGURE 77-16** Myeloid hyperplasia. This marrow shows an increase in the fraction of cells in the myeloid or granulocytic lineage as might be seen in a normal marrow responding to infection. The myeloid/erythroid (M/E) ratio is  $>3:1$ . (From RS Hillman et al: *Hematology in Clinical Practice*, 5th ed. New York, McGraw-Hill, 2010.)